

# REDUCED EXPRESSIONS IN SEMIDIRECT PRODUCTS OF COXETER GROUPS

TOM EDGAR<sup>1</sup>

ABSTRACT. Any normal reflection subgroup  $\widetilde{W}$  of a Coxeter system  $(W, S)$  is a factor in a semidirect product decomposition of  $W$  as described by Bonnafé and Dyer. Namely,  $S$  is the union of two subsets  $I$  and  $J$  such that no element of  $I$  is conjugate to an element of  $J$ ,  $\widetilde{W}$  is the subgroup generated by  $W_I$ -conjugates of elements of  $J$ , and  $W$  is the semidirect product of  $W_I$  by  $\widetilde{W}$ . This note describes the reduced expressions of elements of the form  $wxw^{-1}$  with  $w \in W_I$  and  $x \in W_J$  in terms of reduced expressions of  $x$  and a suitable element of  $W_I$ .

## INTRODUCTION AND PRELIMINARIES

Following Bonnafé and Dyer, we consider a Coxeter system  $(W, S)$  in its internal semidirect product decomposition with respect to two subsets  $I, J \subset S$  such that  $I \cap J = \emptyset$  and  $I \cup J = S$  and with the requirement that no element of  $I$  is conjugate to an element of  $J$ . The latter statement says that the vertex set of the Coxeter graph of  $(W, S)$  can be divided into two subsets with only edges with even label connecting the two. Let  $T$  represent the set of reflections of  $W$ . In [4] (see also [1]), it is shown that  $W$  is the semidirect product of  $W_I$  and the normal subgroup  $\widetilde{W}$  generated by  $W_I$ -conjugates of elements of  $J$ . We observe that any normal reflection subgroup  $\widetilde{W}$  of  $W$  arises from this type of decomposition of  $S$ . Then, we describe the reduced expressions of elements of the form  $wxw^{-1}$  where  $w \in W_I$  and  $x \in W_J$  in terms of reduced expressions of  $w$  and  $x$ . From this, we get a slightly stronger result describing reduced expressions of reflections of the form  $wtw^{-1}$  with  $t \in W_J \cap T$  and  $w \in W_I$ . For general results about Coxeter systems consult [5].

### 1. STATEMENT OF MAIN RESULT

We begin with the same setup as [1]. We let  $(W, S)$  be a Coxeter system with  $\ell : W \rightarrow \mathbb{N}$  the corresponding length function. Furthermore, assume that  $S = I \cup J$  with  $I \cap J = \emptyset$  and that no element of  $I$  is conjugate to an element of  $J$ . We denote by  $W_I$  and  $W_J$  the standard parabolic subgroups generated by  $I$  and  $J$  respectively. For any  $K, L \subset S$ , we let  $W^K := W/W_K$  (respectively  ${}^L W := W_L \backslash W$ ) be the set of shortest right coset representatives of  $W_K$  in  $W$  (respectively the set of shortest left coset representatives of  $W_L$  in  $W$ ). Then we define  ${}^L W^K := {}^L W \cap W^K$  to be the set of shortest representatives for the double cosets  $W_L \backslash W / W_K$  in  $W$ . For brevity, we will use the following notation  $W^K_L := W^K \cap W^L$ . Additionally, let  $T = \cup_{w \in W} wSw^{-1}$  be the set of reflections of  $W$ . If  $\mathcal{P}(T)$  denotes the power set of

---

*Date:* February 20, 2009.

<sup>1</sup>Department of Mathematics, University of Notre Dame, Notre Dame, IN 46556; tedgar@nd.edu

$T$ , then we let  $N : W \rightarrow \mathcal{P}(T)$  be given by  $N(w) = \{t \in T \mid \ell(tw) < \ell(w)\}$ . We define a new set  $\tilde{J} = \{wsw^{-1} \mid w \in W_I; s \in J\} \subset T$ , and then let  $\tilde{W}$  be the subgroup of  $W$  generated by  $\tilde{J}$ . Part a) of the following result is due to [4, Proposition 2.1]; part b), along with a stronger version of part a), is proven (independently) in [1].

**Theorem 1.1** (Gal; Bonnafé, Dyer). *Let  $(\tilde{W}, \tilde{J})$  be as above.*

- a)  $(\tilde{W}, \tilde{J})$  is a Coxeter system and  $W = \tilde{W} \rtimes W_I$ .
- b) Every element  $w \in W_I$  is the unique element of the coset  $w\tilde{W}$  with minimal length.

We complement [1] by noting that any normal reflection subgroup of a Coxeter group appears as the normal component of a semidirect product decomposition for  $W$  as in Theorem 1.1.

**Proposition 1.2.** *Let  $W'$  be any normal reflection subgroup of  $W$ . Then, there exists a subset  $I \subset S$  such that  $W = W_I \rtimes W'$ , that is, any normal reflection subgroup of  $W$  is a normal semidirect factor of  $W$ .*

*Proof.* Let  $W''$  be any reflection subgroup of  $W$  and let  $\tilde{W}'$  be its normal closure in  $W$ . Then, define  $J \subset S$  to be  $J = S \cap (\bigcup_{w \in W} w(W'' \cap T)w^{-1})$ . Since  $W'$  is normal and generated by the reflections it contains, we have that  $W'$  is generated by the  $W$ -conjugates of  $J$ . Now, we let  $I = S \setminus J$ , and let  $\tilde{J} = \{wsw^{-1} \mid w \in W_I; s \in J\}$  and  $\tilde{W}$  be the group generated by  $\tilde{J}$ . By Theorem 1.1, we know that  $\tilde{W}$  is normal; since  $J \subset \tilde{W}$ , we have  $wJw^{-1} \subset \tilde{W}$  for all  $w$  and so  $W' \leq \tilde{W}$ . Thus,  $W' = \tilde{W}$ . Also by the theorem, we know that  $W = W_I \rtimes \tilde{W}$  so  $W = W_I \rtimes W'$ .  $\square$

So any normal reflection subgroup arises from a decomposition of  $S$  into two disjoint subsets such that no element in one set is conjugate to an element of the other set.

Now, we can move on to our main result. For any element  $x \in W$ , define the set

$$x^\perp := \{s \in S \mid sr = rs \text{ for all } r \text{ appearing in a reduced expression for } x\},$$

which is well defined since any reduced expression for  $x$  uses the same set of simple reflections.

Now, for fixed  $w \in W_I$ , elements of the form  $wxw^{-1}$  with  $x \in W_J$  form a standard parabolic subgroup of  $\tilde{W}$ , namely the subgroup generated by the set

$$J' = \{wsw^{-1} \mid s \in J\} \subset \tilde{J}.$$

In Lemma 2.4 we will show that every  $W_I$ -conjugate of an element in  $W_J$  is uniquely expressible as  $wxw^{-1}$  with  $w \in W_I^{I \cap x^\perp}$  and  $x \in W_J$ . Our goal is to understand how reduced expressions of elements of this form reflect the semidirect product decomposition of  $W$ .

For any expression  $r_1 \dots r_n$  with  $r_i \in S$  for all  $i$  and  $D \subset \{1, \dots, n\}$ , we denote by  $\prod_D r_i$  the product  $r_{i_1} \dots r_{i_j}$  where  $i_1 < i_2 < \dots < i_j$  and  $D = \{i_1, \dots, i_j\}$ . The main result here characterizes the reduced expressions of elements of the form  $wxw^{-1}$  where  $w \in W_I$  and  $x \in W_J$ .

**Theorem 1.3.** *Suppose that  $S = I \cup J$ ,  $I \cap J = \emptyset$ , and no element of  $I$  is conjugate to an element of  $J$ , and define  $(\tilde{W}, \tilde{J})$  as above. Suppose that  $x \in W_J$  and  $w \in W_I^{I \cap x^\perp}$  with  $\ell(x) = m$ . Fix a reduced expression  $wxw^{-1} = r_1 \dots r_d$  for  $wxw^{-1}$ . From*

this expression, we can define two sets  $A_J := \{i \mid r_i \in J\}$  and  $A_I := \{i \mid r_i \in I\}$ . Then the following are true:

- (1)  $|A_J| = m$ .
- (2)  $\prod_{A_J} r_i$  is a reduced expression of  $x$ .
- (3) There exists  $B \subset A_I$  where  $\prod_B r_i$  is a reduced expression of  $w$  and  $\prod_{A_I \setminus B} r_i$  is a reduced expression of  $w^{-1}$ .
- (4) Let  $i \in A_J$ , so  $r_i \in J$ . Then
  - (a)  $r_i r_j = r_j r_i$  for all  $j \in B$  with  $j > i$ .
  - (b)  $r_i r_k = r_k r_i$  for all  $k \in A_I \setminus B$  with  $k < i$ .
- (5) Let  $i \in B$  and  $j \in A_I \setminus B$  with  $j < i$ . Then  $r_i r_j = r_j r_i$ .
- (6)  $N(wxw^{-1}) \cap \widetilde{W} = wN(x)w^{-1}$ .

This theorem basically says that every reduced expression for  $wxw^{-1}$  is essentially of the form: (reduced expression of  $w$ )(reduced expression of  $x$ )(reduced expression of  $w^{-1}$ ) up to commutation relations. Following this, we can say a bit more for the case of  $x$  being a reflection  $x = t \in W_J \cap T$ .

**Corollary 1.4.** *Consider the same setup as Theorem 1.3. Let  $w \in W_I^{I \cap x^\perp}$  and  $x = t \in W_J \cap T$  with  $\ell(t) = 2k + 1$  so that  $wtw^{-1} = r_1 \dots r_{e+1} \dots r_{2e+1}$  is a reduced expression. Then following the notation from Theorem 1.3, the following are true:*

- (1)  $e + 1 \in A_J$  and  $|\{i \in A_J \mid i < e + 1\}| = k$ .
- (2)  $B = \{i \in A_I \mid i < e + 1\}$  and so  $A_I \setminus B = \{i \in A_I \mid i > e + 1\}$ .

## 2. PROOF OF THEOREM 1.3

To prove the main result, we need two auxiliary results. To get to these results, we first remind the reader of the following standard results regarding shortest double coset representatives.

**Proposition 2.1.** *Let  $K, L \subset S$ . The following hold:*

- (1) For  $w \in {}^K W^L$ ,  $W_K \cap wW_L w^{-1} = W_{K \cap wLw^{-1}}$ .
- (2) Let  $w \in {}^K W^L$ , then any  $x \in W_K wW_L$  has a unique expression  $x = uwv$  with  $u \in W_K^{K \cap wLw^{-1}}$  and  $v \in W_L$ . Furthermore, for this expression,  $\ell(uvw) = \ell(u) + \ell(w) + \ell(v)$ .

*Proof.* See [2, 2.7]. The proofs given are for finite Coxeter groups, but the proofs easily extend to any Coxeter group.  $\square$

This proposition is the main ingredient for the proofs of Lemma 2.3 and Lemma 2.4 below. To prove these lemmas, it is also useful to have the following:

**Lemma 2.2.** *Let  $I, J \subset S$ ,  $I \cap J = \emptyset$ , and let  $x \in W_J$ . Then  $x \in {}^I W^I$  and  $I \cap xIx^{-1} = I \cap x^\perp$ .*

*Proof.* Clearly multiplying  $x$  by a simple reflection from  $I$  on the left or right must increase the length so  $x \in {}^I W^I$ . Now the fact that  $I \cap xIx^{-1} \subset I \cap x^\perp$  follows since if  $r = xsx^{-1}$  for some  $r, s \in I$ , then  $rx = xs$ . Now, both  $rx$  and  $xs$  are reduced expressions (for any reduced expression of  $x$ ) and since all reduced expressions use the same simple reflections we must have  $r = s$  since  $x \in W_J$ . Thus we have  $r \in x^\perp$ . Containment in the other direction is trivial.  $\square$

**Lemma 2.3.** *Let  $I, J \subset S$  with  $I \cap J = \emptyset$ . Let  $x \in W_J$ , and let  $w \in W_I^{I \cap x^\perp}$  (that is,  $w$  is in  $W_I$  and is a shortest right coset representative of  $W_{I \cap x^\perp}$ ). Then  $\ell(wxw^{-1}) = 2\ell(w) + \ell(x)$ .*

*Proof.* This is just a combination of Proposition 2.1 and Lemma 2.2.  $\square$

The next lemma, which follows from Proposition 2.1, gives a normal form for  $W_I$ -conjugates of reflections in  $W_J$ .

**Lemma 2.4.** *Suppose that  $S = I \cup J$  and  $I \cap J = \emptyset$ . Every  $W_I$ -conjugate of an element in  $W_J$ ,  $wxw^{-1}$  ( $w \in W_I$ ,  $x \in W_J$ ) is uniquely expressible as  $uxu^{-1}$  with  $u \in W_I^{I \cap x^\perp}$  and  $x \in W_J$ .*

*Proof.* Since  $x \in {}^I W^I$  then by Proposition 2.1 part (2), we know that  $wxw^{-1} = uxv$  with  $u \in W_I^{I \cap x^\perp}$  (by Lemma 2.2) and  $v \in W_I$ . We note that  $w = u'w'$  where  $u' \in W_I^{I \cap x^\perp}$  and  $w' \in W_{I \cap x^\perp}$  so that  $wxw^{-1} = u'w'xw^{-1} = u'xw'w^{-1} = u'xu'^{-1}$  so that by necessity  $u' = u$  and so  $v = u^{-1}$  in the unique decomposition of  $wxw^{-1}$ . Thus,  $wxw^{-1} = uxu^{-1}$  with  $u \in W_I^{I \cap x^\perp}$ .  $\square$

*Remark 2.5.* Lemma 2.4 can also be interpreted in the following way. Let  $x, x' \in W_J$  and  $w, w' \in W_I$  then  $wxw^{-1} = w'x'w'^{-1}$  if and only if  $x = x'$  and  $w^{-1}w' \in W_{I \cap x^\perp}$ .

From this point on, we will assume that  $S = I \cup J$  with  $I \cap J = \emptyset$  and no element of  $I$  is conjugate to an element of  $J$ .

*Remark 2.6.* Before we move the proof of Theorem 1.3, notice that since  $\widetilde{W} \triangleleft W$ , then for  $w$  and  $x$  as in Theorem 1.3 with  $wxw^{-1} = r_1 \dots r_d$  we have  $t_i := r_1 \dots r_{i-1} r_i r_{i-1} \dots r_1 \in \widetilde{W}$  if and only if  $r_i \in J$ .

*Proof of Theorem 1.3.* Suppose that  $w = u_1 \dots u_n$  and  $x = s_1 \dots s_m$  are reduced expressions for  $w$  and  $x$  respectively. Then, we know by Lemma 2.3 that  $wxw^{-1} = u_1 \dots u_n s_1 \dots s_m u_n \dots u_1$  is a reduced expression for  $wxw^{-1}$ . Therefore, according to Remark 2.6,  $|N(wxw^{-1}) \cap \widetilde{W}| = \ell(x)$ . Using the remark in reverse, if  $wxw^{-1} = r_1 \dots r_{2n+m}$  is any reduced expression, there must be  $i_1 < i_2 < \dots < i_k$  such that  $r_{i_j} \in J$  for all  $j$  and  $k = \ell(x)$ . Then  $A_J = \{i_1, \dots, i_{\ell(x)}\}$ , and this proves (1).

Now, any reduced expression can be achieved by applying braid operations to a given reduced expression. We already have a particular reduced expression, namely  $wxw^{-1} = u_1 \dots u_n s_1 \dots s_m u_n \dots u_1$ , so  $A_J = \{n+1, \dots, n+m\}$ ,  $A_I = \{1, \dots, n, n+m+1, \dots, 2n+m\}$  and  $B = \{1, \dots, n\}$ . Thus (2)-(5) clearly hold for this reduced expression. Now we proceed to prove (2)-(5) by induction on the number of braid operations needed to change from  $u_1 \dots u_n s_1 \dots s_m u_n \dots u_1$ .

Suppose  $wxw^{-1} = r_1 \dots r_{2n+m}$  is a reduced expression satisfying (2)-(5) of the theorem. For this expression we have  $A_J, A_I$  and  $B$  defined. Clearly, braid operations involving simple reflections,  $r_i$ , with indices,  $i$ , that are both in  $A_J$ , both in  $B$ , or both in  $A_I \setminus B$  are just rearranging  $x, w$ , or  $w^{-1}$  respectively and are thus giving new reduced expressions for  $x, w$ , or  $w^{-1}$  respectively; consequently, the theorem holds. This leaves us with three cases.

**Case 1:**  $r_i \in I$  and  $r_{i+1} \in J$  (or vice versa). If  $r_i$  and  $r_{i+1}$  do not commute, then we know that  $r_{i-1} = r_{i+1} \in J$  or  $r_{i+3} = r_{i+1} \in J$  by the restriction that no element of  $I$  is conjugate to an element of  $J$ . However, either of these situations contradicts the fact that  $\prod_{A_J} r_i = r_{i_1} \dots r_{i_{\ell(x)-1} r_{i_{\ell(x)}} r_{i_{\ell(x)-2}} \dots r_{i_1}$  is a reduced expression of  $x$ , which holds because  $wxw^{-1} = r_1 \dots r_{2n+m}$  is

a reduced expression satisfying (2). Therefore,  $r_i$  commutes with  $r_{i+1}$  and thus  $wxw^{-1} = r_1 \dots r_{i+1} r_i \dots r_{2n+m}$  is a reduced expression satisfying (2)-(5).

**Case 2:**  $i \in B$  and  $i+1 \in A_I \setminus B$  so  $r_i \in I$  and  $r_{i+1} \in I$ . Let  $r := r_i$  and  $s := r_{i+1}$  and suppose that these simple reflections do not commute. Then to perform a braid operation, either  $r_{i-1} = s$  or  $r_{i+2} = r$  (or both). Without loss of generality, we assume  $r_{i+2} = r$ . We note that in this case we must have  $i+2 \in A_I \setminus B$  otherwise we contradict the fact that  $\prod_B r_i$  is a reduced expression for  $w$ . However, by induction, we know that  $r = r_i$  commutes with all  $r_j$  with  $j < i$  and  $j \in A_J$  and simultaneously,  $r = r_{i+2}$  commutes with all  $r_j$  with  $j > i+2$  and  $j \in A_J$ . This implies that  $r$  commutes with all  $r_j$  with  $j \in A_J$  so that  $r$  commutes with  $x$ . Finally, we see also by induction that  $r = r_{i+2}$  commutes with all  $r_k$  with  $k \in B$  and  $k > i$ . Thus,  $w$  has a reduced expression ending in  $r$  and  $r$  commutes with  $x$  contradicting the fact that  $w \in W_I^{I \cap x^\perp}$ . So,  $r_i$  must commute with  $r_{i+1}$  and  $wxw^{-1} = r_1 \dots r_i r_{i+1} r_{i+2} \dots r_{2n+m}$  is a new reduced expression satisfying (2)-(5).

**Case 3:**  $i \in A \setminus B$  and  $i+1 \in B$  so  $r_i \in I$  and  $r_{i+1} \in I$ . Since the reduced expression we are currently in satisfies the theorem by induction, property (5) implies that  $r_i$  and  $r_{i+1}$  commute.

So we have shown that any braid operation that can occur must either be a commutation relation or be a rearrangement of  $w$ ,  $x$ , or  $w^{-1}$  separately.

Lastly, we prove part (6). As we have seen,  $|N(wxw^{-1}) \cap \widetilde{W}| = \ell(x) = |N(x)|$ . Clearly,  $wN(x)w^{-1} \subset \widetilde{W}$  since  $x \in W_J$ ,  $W_J \subset \widetilde{W}$  and  $\widetilde{W}$  is normal. So let  $t \in N(x)$ , that is  $\ell(tx) < \ell(x)$ . Since  $\ell(wxw^{-1}) = 2\ell(w) + \ell(x)$ , then  $\ell(wtw^{-1}wxw) = \ell(wtxw^{-1}) < \ell(wxw^{-1})$  because  $tx$  has smaller length than  $x$ . Thus,  $wtw^{-1} \in N(wxw^{-1})$  for all  $t \in N(x)$ , and so  $wN(x)w^{-1} \subset N(wxw^{-1})$ . Finally, we have seen that  $wN(x)w^{-1} \subset N(wxw^{-1}) \cap \widetilde{W}$  and these sets are both finite with the same cardinality so they must be equal.  $\square$

Lastly, the corollary follows from the following information about reflections. For a reflection  $t$ , we say that  $t = r_1 r_2 \dots r_n$  is a palindromic reduced expression if  $r_i = r_{n-i+1}$  for all  $i \leq \frac{n-1}{2}$ . The following lemma is due to [3].

**Lemma 2.7.** *If  $t = r_1 \dots r_{2m+1}$  is a reduced expression for  $t \in T$ , then  $t = r_1 \dots r_m r_{m+1} r_m \dots r_1$  is a reduced palindromic expression of  $t$ .*

**Corollary 2.8.** *Suppose  $I$  and  $J$  are disjoint,  $S = I \cup J$  and no element of  $I$  is conjugate to any element of  $J$ . Let  $w \in W_I$  and  $t \in W_J \cap T$ . Now, for any reduced expression  $wtw^{-1} = r_1 \dots r_{n+1} \dots r_{2n+1}$  then we have  $wtw^{-1} = r_1 \dots r_n r_{n+1} r_n \dots r_1$  and so  $r_{n+1} \in J$ .*

*Proof.* This is just a restatement of Lemma 2.7 along with the fact that  $wtw^{-1} \in \widetilde{W}$  so that  $wtw^{-1}$  must be conjugate to an element of  $J$ .  $\square$

*Proof of Corollary 1.4.* Let  $wtw^{-1} = r_1 \dots r_{e+1} \dots r_{2e+1}$ . The proof follows directly from Corollary 2.8, which tells us that  $e+1 \in A_J$ . Thus, we can never have a braid operation occur between the middle element and a simple reflection in  $I$ . Therefore, we can never have  $i \in B$  and  $j \in A_I \setminus B$  with  $j < i$  since that would have required a braid relation between  $r_i$  and the middle element or  $r_j$  and the middle element, which would in turn have given us a reduced expression for  $wtw^{-1}$  with  $r_{e+1} \in I$ .  $\square$

## REFERENCES

- [1] C. Bonnafé and M.J. Dyer. Semidirect product decomposition of Coxeter groups. preprint (2008), available at [arXiv:0805.4100](https://arxiv.org/abs/0805.4100).
- [2] Roger W. Carter. *Finite groups of Lie type*. Pure and Applied Mathematics (New York). John Wiley & Sons Inc., New York, 1985. Conjugacy classes and complex characters, A Wiley-Interscience Publication.
- [3] Matthew Dyer. Reflection subgroups of Coxeter systems. *J. Algebra*, 135(1):57–73, 1990.
- [4] Światosław R. Gal. On normal subgroups of Coxeter groups generated by standard parabolic subgroups. *Geom. Dedicata*, 115:65–78, 2005.
- [5] James E. Humphreys. *Reflection groups and Coxeter groups*, volume 29 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1990.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NOTRE DAME, NOTRE DAME, IN 46556  
E-mail address: [tedgar@nd.edu](mailto:tedgar@nd.edu)